

## Application of Gauss - Lanczos Algorithm to Determine Low Modes Density of Dirac Operator

Rudina Osmanaj<sup>a1</sup>, Dafina Xhako<sup>\*b2</sup>, Niko Hyka<sup>c3</sup>

<sup>1</sup>Department of Physics, Faculty of Natural Sciences, University of Tirana, Bvd: "Zog I", Tirana, Albania

<sup>2</sup>Department of Physics Engineering, Faculty of Mathematical Engineering and Physical Engineering, Polytechnic University of Tirana, Tirana, Albania

<sup>3</sup>Department of Diagnostics, Faculty of Medical Technical Sciences, Medical University of Tirana, Tirana, Albania

<sup>a</sup>rudina.zeqirllari@fshn.edu.al; <sup>\*b</sup>dafinaxhako@yahoo.com; <sup>c</sup>nikohyka@gmail.com

### ABSTRACT

There are numerous applications in physics, especially in Lattice QCD, where is required to bound entries and the trace of the inverse and the determinant of a large sparse matrix. This paper review one of the most popular methods which are used in lattice QCD to compute the determinant of the lattice Dirac operator: Gaussian integral representation. A modified algorithm can be used for other purposes too, for example for the determination of the density of eigenvalues of the Dirac operator near the origin. This because in Lattice QCD, low-lying Dirac modes are a suitable tool to understand chiral symmetry since they encode the nature of quark propagation as well as the condensate itself in the chiral regime. The formation of a non-zero chiral condensate is an effect of the accumulation of the low modes of the Dirac operator near zero. We review the development in Krylov subspace evaluation of matrix functions and we develop a practical numerical algorithm to achieve a reliable determination of the density of eigenvalues of the Dirac operator near the origin using the Gauss-Lanczos quadrature. We utilize the optimal properties of Krylov subspaces in approximating the distribution of the eigenvalues of the Dirac operator. In this work we have used the Boriçi - Creutz operator to test our method, as an example of using this algorithm in Lattice QCD.

**Keywords:** algorithm; Boriçi - Creutz operator; eigenvalues; Dirac operator; Gauss-Lanczos quadrature; lattice QCD; numerical simulation.

### 1. INTRODUCTION

Lattice QCD is a well - established non - perturbative approach to solving the quantum chromodynamics (QCD) theory of quarks and gluons. It is a lattice gauge theory formulated on a grid or lattice of points in space and time. When the size of the lattice is taken infinitely large and its sites infinitesimally close to each other, the continuum QCD is recovered [20]. But the simulation of light fermions on a lattice is always a challenging task. There are several prescriptions to circumvent or minimize the doubling problem without spoiling the chiral symmetry on the lattice. By the no-go theorem [16], the

minimum number of species one can have on a lattice with chiral symmetry is two what is known as a "minimally doubled" fermion.

There are several formulations of minimally doubled fermions, but we will be focused on Boriçi - Creutz (BC) fermions [4], [6]. These fermions share the desirable features of strict locality with traditional discretization, while preserving exact chiral symmetry for a degenerate doublet of quark fields, but breaks the hypercubic symmetry [3]. In order to correct this broken symmetry, can be add non-covariant counter terms through quantum corrections. The renormalization properties of the BC fermions at one loop in the perturbation theory have been investigated in reference [5], [21] and the free Dirac operator for Boriçi - Creutz fermions in momentum space is written as:

$$D_{BC}(p) = \sum_{\mu} \left[ i\gamma_{\mu} \sin(p_{\mu}) + i(\Gamma - \gamma_{\mu}) \cos(p_{\mu}) \right] + i(c_3 - 2)\Gamma, \quad (1)$$

where  $c_3$  is one of the counter terms added in the action by the authors of ref [5], [21], the most relevant term that contributes in the corrected action. The aim of the original study is to restore the broken symmetry by studying the chiral condensate, but in this paper, we will focus on the algorithm we have used in order to perform a part of the study, the Gauss - Lanczos quadrature. [1]

## 2. THE EFFECTIVE CONDENSATE

The Banks - Casher relation [2] links the spectral density  $\rho(\lambda, m)$  of the Dirac operator to the condensate as:

$$\lim_{\lambda \rightarrow 0} \lim_{m \rightarrow 0} \lim_{V \rightarrow \infty} \rho(\lambda, m) = \frac{\Sigma}{\pi}, \quad (2)$$

where

$$\Sigma = -\lim_{m \rightarrow 0} \lim_{V \rightarrow \infty} \langle \bar{\psi} \psi \rangle. \quad (3)$$

Let remember that in a space - time box of volume  $V$  with periodic or anti-periodic boundary conditions, the Euclidian massless Dirac operator  $D$  in presence of a given gauge field has purely imaginary eigenvalues  $i\lambda_1, i\lambda_2, \dots$ , which may be ordered so that those with the lower magnitude come first. The associated average spectral density is given by:

$$\rho(\lambda, m) = \frac{1}{V} \sum_{k=1}^{\infty} \langle \delta(\lambda - \lambda_k) \rangle, \quad (4)$$

where the bracket  $\langle \dots \rangle$  denotes the QCD (Quantum Chromo Dynamics) expectation value and  $m$  the current - quark mass.

The Banks - Casher relation can be read in both directions: a non-zero spectral density implies that the symmetry is broken by a non - vanishing  $\Sigma$  and vice - versa. As we mentioned above, spectral density and low - lying Dirac modes are very important for understanding spontaneous chiral symmetry breaking. Furthermore, the chiral condensate is a useful order parameter to distinguish the phases at  $T = 0$ , for minimally doubled fermions [4], [7], [12], [20], [23-27]. In the continuum theory and for small masses, the mode number can be calculated analytically in chiral perturbation theory. In infinite

volume, chiral perturbation theory yields an expansion of the spectral density  $\rho(\lambda, m)$  essentially in powers of  $\lambda$  and  $m$ . The leading term is given by the Banks - Casher formula which relates the mode number to the "effective chiral condensate" defined through:

$$\Sigma_{eff} = \frac{\pi}{2} \frac{\nu(\Lambda)}{\Lambda V}, \quad (5)$$

where

$$\Lambda = \sqrt{M^2 - m^2}. \quad (6)$$

So, instead of using spectral density, the average number  $\nu(\Lambda)$  of eigenmodes of the massive Hermitian operator  $D^\dagger D + m^2$  with eigenvalues  $\alpha \leq M^2$ , where  $M^2$  is a chosen cutoff value, seems to be a more convenient quantity to deal with. Since:

$$\nu(\Lambda) = V \int_{-\Lambda}^{\Lambda} \rho(\lambda, m) d\lambda, \quad (7)$$

the mode number carries the same information as the spectral density [8], [10], [18], [23-27]. In the presence of a given gauge field, the number of eigen-modes of with eigenvalues can be determined by calculating the eigenvalues and their multiplicities numerically.

Spectral density and low-lying Dirac modes calculation is very important in order to understand the dynamics of spontaneous chiral symmetry breaking. There are several articles and studies on the spectral density of the Dirac operator, such as in references [6], [9], [10]. In these papers the Dirac modes number is calculated by approximating the operator for low-lying modes applying the spectral projector method. While in our work we propose a spectral method for the determination of the spectral density of Dirac operator near the origin using Gauss - Lanczos quadrature and we can show the dynamical spontaneous chiral symmetry breaking using chiral lattice fermions (Boriçi - Creutz fermions).

### 3. GAUSS - LANCZOS ALGORITHM

In this section, we show an alternative way to count efficiently the mode number and estimate the effective condensate. We utilise the optimal properties of Krylov subspaces in approximating the distribution of the eigenvalues of the Dirac operator. These properties allow us to use the Gauss - Lanczos quadrature in order to estimate the number of modes of the Dirac operator. What we propose is to use Lanczos algorithm in order to produce the tridiagonal and symmetric Lanczos matrix  $T_n$ . Its eigenvalues, or the so-called Ritz values, tend to approximate the eigenvalues of the original matrix  $A_n = D_n^\dagger D_n$ . The eigenvalues and eigenvectors of a symmetric and tridiagonal matrix can be computed by the QR method [1], [11]. In fact, Ritz values are the nodes of the Gauss - Lanczos quadrature with the corresponding weights being the first elements of the Ritz vectors squared. Bai, Fahey and Golub [1] observed that a vector-matrix-vector of the form  $\psi^\dagger f(A) \psi$  can be expressed as an integral of the function  $f(\cdot)$  over a modified spectral measure. An  $n$ -point Gaussian quadrature approximation to this integral is then given by

$$\psi^\dagger f(A) \psi \approx \sum_{i=1}^n \omega_i f(\theta_i), \quad (8)$$

where the abscissas,  $\theta_i$ , of this quadrature rule are given by the Ritz values. These are generated by the Lanczos algorithm applied with seed vector  $\psi$ , while the weights  $\omega_i$  are the squares of the first components of the corresponding eigenvectors of  $T_n$ .

So, using Gauss - Lanczos quadrature we calculate the distribution of the eigenvalue's modes, and then we find the low-lying Dirac modes as:

$$\frac{\nu(\Lambda)}{V} = \sum_{i=1}^{k(\Lambda)} \omega(\theta_i), \quad (9)$$

since our function  $f(\theta_i)$  is the Heaviside function:

$$f(\theta_i) = \begin{cases} 1 & i \leq k(\Lambda) \\ 0 & i > k(\Lambda) \end{cases}. \quad (10)$$

Let  $A \in \mathbb{C}^{N \times N}$  be a Hermitian matrix and  $b \in \mathbb{C}^N$  a starting vector. Then the following algorithm computes the Gauss - Lanczos quadrature.

#### Gauss - Lanczos algorithm

---

Calculate  $\alpha_i$  dhe  $\beta_i$  using Lanczos algorithm for  $Ax=b$   
 Define  $(T_n)_{i,i} = \alpha_i$ ,  $(T_n)_{i+1,i} = (T_n)_{i,i+1} = \beta_i$ ,  $(T_n)_{i,j} = 0$   
 Calculate the eigenvalue and eigenvectors of the matrix  $T_n$ , where  $i=1 \dots n$   
 Sort the eigenvalues and eigenvectors in the increasing order of eigenvalues  
 Define  $k$  as the maximal index  $s$ , the maximum index which correspond to the cut-off eigenvalue  
 Define  $\theta_i$  as the positive square root of the original eigenvalue  
 Define  $z_i$  as the first elements of the eigenvectors  $v_i$ ,  $i = 1 \dots n$   
 Define  $\omega_i = z_i^2$   
 Calculate the mode number  $\mathcal{G}_k = \sum_i^k \omega_i$

---

Let remember that the Lanczos algorithm (C. Lanczos, 1952) is as below: Let be  $D$  the Dirac matrix. In many cases in order to find the eigenvalues of a huge matrix, is sufficient to find the eigenvalues of the respective tridiagonal matrix, which eigenvalues are very close to those of the original matrix. Using Lanczos algorithm, we define the element of the tridiagonal matrix.

#### Lanczos algorithm

---

% Let be  $A = D^*D$  and  $b = z_2$   
 %  $\beta_0 = 0, \rho_1 = \|b\|_2^{-1}, q_0 = 0, q_1 = \rho_1 b$   
**for**  $i = 1 \dots n$  **do**  
      $v = Aq_i$   
      $\alpha_i = q_i^* v$   
      $v := v - q_i \alpha_i - q_{i-1} \beta_{i-1}$   
      $\beta_i = \|v\|_2$   
      $\rho_{i+1} = \frac{v}{\beta_i}$

$$\rho_{i+1} = -\frac{(\rho_i \alpha_i + \rho_{i-1} \beta_{i-1})}{\beta_i}$$

if  $\left| \frac{\rho_i}{\rho_{i+1}} \right| < tol$  then  
 $n = i$   
 stop  
 end if  
 end for

#### 4. RESULTS

In order to test our method, we have used BC fermions in the quenched approximation for  $\beta = 6$  in a  $24^4$  lattice. For this kind of lattice fermions, the flavor number is  $n_f = 2$  and the counter term we have chosen is  $c_3 = 0.4$ . Using Gauss - Lanczos quadrature we have calculated the mode number for minimally doubled fermions BC, and therefore the effective chiral condensate, in a background of 1000 gauge configurations, generated using the SU(3) theory. The simulations are made using the Hybrid Monte Carlo (HMC) algorithm and for the calculations we have used Lanczos algorithm [13], implemented in FermiQCD, a collection of C++ classes, functions and parallel algorithms for lattice QCD, based on Matrix Distributed Processing [22]. As a condition in order to stop the Lanczos algorithm we have set that the residue is smaller than the tolerance  $tol = 10^{-3}$ , because the number of modes of the Dirac operator for 1000 configuration, considering the errors too, doesn't change for different values of the tolerance. This can be seen in the Figure 1, where is presented the number of modes for different tolerances for the minimally doubled fermions Boriçi – Creutz. The results are averaged over 1000 configurations. As expected by the chiral perturbation theory, the low modes of the operator condense and reach a "plateau", as can be seen in the Figure 2.




Figure 1. In this figure is presented the number of modes for certain eigenvalues for different tolerances for the minimally doubled fermions Boriçi – Creutz.




Figure 2. In this figure can be seen clearly the condensation of low modes of the Dirac Operator and the formation of a non-zero effective chiral condensate.

## 5. CONCLUSIONS

Although In this paper we presented one of the uses of Gauss - Lanczos quadrature in Lattice QCD: the evaluation of the low - lying Dirac modes. The methodology we propose, utilize the optimal properties of Krylov subspaces in approximating the distribution of the eigenvalues of the Dirac operator, not in calculating every eigenvalue of the operator. This is distinct from an earlier method where the integrated spectral density (mode number) was calculated efficiently for some preselected fixed range of integration. This algorithm can be used in Lattice QCD for different scopes, for exploring the phase structure of QCD for example, as was the main aim of our original study. Using this algorithm and the Boriçi - Creutz action with zero quark mass in the quenched approximation we take results that give an "expected" qualitative "behavior" of low-lying modes as predicted by the chiral theory.

## ACKNOWLEDGMENT

The authors wish to thank the Department of Physics, Faculty of Natural Science and the Department of Engineering Physics of Polytechnic University of Tirana, Albania.

## REFERENCES

- [1] Fahey G Bai Zh and Golub G. Some large-scale matrix computation problem. In: Journal of Computational and Applied Mathematics. vol. 741. no. 2. 1996. pp. 71–89.
- [2] Banks T and Casher A. Chiral symmetry breaking in confining theories. In: Nuclear Physics B. vol. 169. no.1. 1980. pp. 103–125.
- [3] Bedaque et al. P. Broken symmetries from minimally doubled fermions. In: Physics Letters B, vol. 662, no. 5, 2008, pp. 449–455.

- [4] Boriçi A. Minimally Doubled Fermion Revival. In: PoS LATTICE2008, 2008, p. 231.
- [5] Capitani et al. S. Renormalization of minimally doubled fermions. In: Journal of High Energy Physics, vol.9, 2010, pp. 27-32.
- [6] Creutz M. Four-dimensional graphene and chiral fermions. In: Journal of High Energy Physics, vol.4, 2008, pp. 017-022.
- [7] Damgaard et al. P H. The microscopic spectral density of the QCD Dirac operator. In: Nuclear Physics B, vol. 547, no.1, 1999, pp 305–328.
- [8] Fukaya et al. H. Determination of the chiral condensate from QCD Dirac spectrum on the lattice. In: Physical review D, vol. 83, no. 7, 2011, pp. 074501.
- [9] Giusti L and Lüscher M. Chiral symmetry breaking and the Banks-Casher relation in lattice QCD with Wilson quarks. In: Journal of High Energy Physics, vol.03, 2009, p. 013.
- [10] Golub G and Van Loan C. Matrix computations. In: JHU Press. 2013. pp. 105-153.
- [11] Karsten L. Lattice fermions in Euclidean space-time. In: Physics Letters B. vol. 104. no.4. 1981. pp. 315–319.
- [12] Kogut J and Susskind L. Hamiltonian formulation of Wilson’s lattice gauge theories. In: Physical Review D. vol. 11. no. 2. 1975. pp. 395-342.
- [13] Lanczos C. Solution of systems of linear equations by minimized iterations. In: J. Res. Nat. Bur. Standards. vol. 49. no.1. 1952. pp. 33–53.
- [14] Nambu Y and Jona-Lasinio G. Dynamical model of elementary particles based on an analogy with superconductivity. In: Physical Review. vol. 122. no. 1. 1961. pp. 345-354.
- [15] Neuberger H. Overlap lattice Dirac operator and dynamical fermions. In: Physical Review D. vol. 60. no.6. 1999. p. 065006.
- [16] Nielsen H. and Ninomiya M. Absence of neutrinos on a lattice: (I). Proof by homotopy theory. In: Nuclear Physics B. vol. 185. no. 1. 1981. pp. 20–40.
- [17] Osborn J. Toublan D. Verbaarschot M. From chiral random matrix theory to chiral perturbation theory. In: Nuclear Physics B. vol. 540. no.1. 1999. pp. 317–344.
- [18] Smilga A and Stern J. On the spectral density of Euclidean Dirac operator in QCD. In: Physics Letters B. vol. 318. no.3. 1993. pp. 531–536.
- [19] Wilczek F. Lattice fermions. In: Physical review letters, vol. 59. no.21. p. 2397. 1987.
- [20] Wilson K. Confinement of quarks. In: Physical Review D. vol. 10. no.8. 1974. p. 2445.
- [21] Capitan S. Creutz M. Weber J. Wittig H. Numerical studies of Minimally Doubled Fermions. In: Journal of High Energy Physics. vol. 10. no.9. 2010. p.27.
- [22] Di Pierro M. Matrix Distributed Processing and FermiQCD. In: Journal of High Energy Physics. 2000. hep-lat/0011083.
- [23] Parman S. and Machmudah A. Waveform based Inverse Kinematics Algorithm of Kinematically Redundant 3- DOF In: International Journal of Innovative Technology and Interdisciplinary Sciences. ISSN: 2613 – 7305. DOI: <https://doi.org/10.15157/IJITIS.2020.3.2.407-428>. Volume 3. Issue 2. 2020. pp. 407- 428.

- [24] Hyka-Xhako D. Osmanaj R. Jani J. The physical scale from string tension in LQCD The 10th Jubilee Conference of the Balkan Physical Union. BPU10 – August 2018. Bulgaria. Proceedings in AIP Conference Proceedings 2075. <https://doi.org/10.1063/1.5091253>. eISSN: 1551-7616. 2019. 110002.
- [25] Xhako D and Zeqirllari R. Chiral fermions algorithms in lattice QCD. In: East European Journal of Physics (East. Eur. J. Phys). DOI:10.26565/2312-4334-2019-1-02. Volume 1. 2019. 34-39.
- [26] Zeqirllari R. and Xhako D. Minimally doubled fermions and spontaneous chiral symmetry breaking. In: EPJ (European Physics Journal) Web of Conferences 175. <https://doi.org/10.1051/epjconf/201817504002>. 2018 pp.04002.
- [27] Zeqirllari R. and Xhako D. Preliminary results of the hypercubic symmetry restauration of Boriçi – Creutz fermions in the light hadrons spectrum. In: AIP (American Institute of Physics) Conference Proceedings 2075. <https://doi.org/10.1063/1.5091257>. 2019. pp. 110006.